

Pseudocritical temperature and equation of state from $N_f = 2$ twisted mass lattice QCD

Florian Burger

Humboldt-Universität zu Berlin

St. Goar – March, 2013



tmfT-Collaboration:

E. M. Ilgenfritz (Joint Institute for Nuclear Research, Dubna)

M. Kirchner, M. Müller-Preußker (HU Berlin),

M. P. Lombardo (INFN Frascati),

C. Urbach (Uni Bonn),

O. Philipsen, L. Zeidlewicz, C. Pinke (Uni Frankfurt)

Outline

Motivation and Setup

T_c and Chiral Scenarios

Thermodynamic Equation of State

Conclusions

Outline

Motivation and Setup

T_c and Chiral Scenarios

Thermodynamic Equation of State

Conclusions

Motivation

- ▶ Explore finite T phase transition/crossover for $N_f = 2$ QCD at vanishing chemical potential
- ▶ Order of transition in the chiral limit not known yet for $N_f = 2$
- ▶ Argued to be in universality class of $O(4)$ 3-dim spin model [R. D. Pisarski, F. Wilczek, 84]
However 1st order not ruled out
(e. g. C. Bonati et al. 2009, C. Bonati et al. 2011)
- ▶ $N_f = 2$ equation of state \rightarrow useful to study onset of mass thresholds in thermodynamics
- ▶ Alternative discretization worthwhile to study systematics
- ▶ Considered observables (so far):
Susceptibilities, renorm. L and $\langle \bar{\psi}\psi \rangle$, equation of state, gluon and ghost propagators (see A. Sternbeck's talk)

Twisted Mass Lattice Regularization

- ▶ $N_f = 2$ Wilson twisted mass action at maximal twist:

$$S_f[U, \psi, \bar{\psi}] = \sum_x \bar{\chi}(x) (1 - \kappa H[U] + 2i\kappa a\mu\gamma_5\tau^3) \chi(x)$$
$$\psi = \frac{1}{\sqrt{2}}(1 + i\gamma_5\tau^3)\chi \quad \text{and} \quad \bar{\psi} = \bar{\chi}\frac{1}{\sqrt{2}}(1 + i\gamma_5\tau^3)$$

- ▶ Advantage: when κ tuned to critical value κ_c :


Automatic $\mathcal{O}(a)$ improvement

[R. Frezzotti, G.C. Rossi, 2004]

- ▶ Disadvantage: explicit flavor symmetry breaking (mostly small, but has to be checked)
- ▶ Tree level improved gauge action:

$$S_g[U] = \beta \left(c_0 \sum_P [1 - \frac{1}{3} \text{ReTr}(U_P)] + c_1 \sum_R [1 - \frac{1}{3} \text{ReTr}(U_R)] \right)$$

Simulation Points



$$T = \frac{1}{N_\tau a}$$

Outline

Motivation and Setup

T_c and Chiral Scenarios


Thermodynamic Equation of State

Conclusions

Signals for the Crossover: Polyakov Loop

- Polyakov loop susceptibility (no convincing signal)


$m_\pi \approx 400$ MeV:




Disconnected Chiral Susceptibility

$$\sigma_{\langle\bar{\psi}\psi\rangle} = V/T (\langle\bar{\psi}\psi^2\rangle - \langle\bar{\psi}\psi\rangle^2)$$


$m_\pi \approx 320$ MeV:



$m_\pi \approx 400$ MeV:




$m_\pi \approx 480$ MeV:




Renormalized $\text{Re}(L)$, $\langle\bar{\psi}\psi\rangle$

$$R_{\langle\bar{\psi}\psi\rangle} = \frac{\langle\bar{\psi}\psi\rangle(T, \mu) - \langle\bar{\psi}\psi\rangle(0, \mu) + \langle\bar{\psi}\psi\rangle(0, 0)}{\langle\bar{\psi}\psi\rangle(0, 0)}$$




$$\langle\text{Re}(L)\rangle_R = \text{Re}(L) \exp(V(r_0)/2T)$$




Renormalized $\text{Re}(L)$, $\langle \bar{\psi}\psi \rangle$

$$R_{\langle \bar{\psi}\psi \rangle} = \frac{\langle \bar{\psi}\psi \rangle(T, \mu) - \langle \bar{\psi}\psi \rangle(0, \mu) + \langle \bar{\psi}\psi \rangle(0, 0)}{\langle \bar{\psi}\psi \rangle(0, 0)}$$



$$\langle \text{Re}(L) \rangle_R = \text{Re}(L) \exp(V(r_0)/2T)$$



Results

Ensemble	B12	C12	D8
$T_c[\text{MeV}], \sigma_{\langle \bar{\psi}\psi \rangle}^2:$	217(5)	229(5)	251(10)
$T_c[\text{MeV}], \langle \text{Re}(L) \rangle_R:$	249(5)	258(5)	266(11)

Signals for deconfinement and restoration of chiral symmetry at different locations in crossover region

$O(4)$ Magnetic EoS

- ▶ Near critical point of a 2^{nd} order transition: universality
- ▶ Expect data to collapse on universal curve [J. Engels & T. Mendes, 99]
- ▶ $\langle \bar{\psi}\psi \rangle = h^{1/\delta} c f(d z) + \underbrace{a_\tau \tau h + b_1 h + \dots}_{\text{scaling violating terms}}$ [S. Ejiri *et al.*, 2009]


$$\tau = \beta - \beta_{\text{chiral}}, \quad h = 2 a \mu, \quad z = \tau/h^{1/(\tilde{\beta}\delta)}$$

- ▶ Connection to spin model:


$$\langle \bar{\psi}\psi \rangle \sim M \text{ (magnetization)}$$

$$\mu \sim H \text{ (external field)}$$

$$\beta \sim T \text{ (temperature)}$$



$O(4)$ Magnetic EoS




Result: $\beta_{\text{chiral}} \approx 3.76(2) \rightarrow$
 $T_c(m_\pi = 0) \approx 166 \text{ MeV}$

Pion Mass:

C12: 480 MeV

B12: 400 MeV


A12: 320 MeV



Chiral Limit Scenarios from Spin Models

- From universal function $h^{1/\delta} f(z)$ one can show:

$$T_c(m_\pi) = T_c(0) + Am_\pi^{2/(\tilde{\beta}\delta)}$$



Outline

Motivation and Setup

T_c and Chiral Scenarios

Thermodynamic Equation of State

Conclusions

Integral Method

▶
$$\epsilon = \left. \frac{T}{V} \frac{\partial \ln Z}{\partial \ln T} \right|_V \quad p = T \left. \frac{\partial \ln Z}{\partial V} \right|_T$$

But: $V = N_\sigma^3 a^3$
 $T = \frac{1}{N_\tau a}$

- ▶ Trace anomaly (interaction measure):

$$I = \epsilon - 3p = - \frac{T}{V} \frac{d \ln Z}{d \ln a}$$

- ▶ Starting point for $p(T)$ and $\epsilon(T)$ via

$$\frac{I}{T^4} = T \frac{\partial}{\partial T} \left(\frac{p}{T^4} \right) \quad \frac{p}{T^4} - \frac{p_0}{T_0^4} = \int_{T_0}^T d\tau \frac{\epsilon - 3p}{\tau^5} \Big|_{\text{LCP}}$$

on lines of constant physics (LCP)

Trace Anomaly




$$\begin{aligned}\frac{l}{T^4} &= \frac{\epsilon - 3p}{T^4} = -\frac{T}{T^4 V} \frac{d \ln Z}{d \ln a} \\ &= N_\tau^4 \left(a \frac{d\beta}{da} \right) \left(\frac{c_0}{3} \langle \text{ReTr} U_P \rangle_{\text{sub}} + \frac{c_1}{3} \langle \text{ReTr} U_R \rangle_{\text{sub}} \right. \\ &\quad \left. + \frac{\partial \kappa_c}{\partial \beta} \langle \bar{\chi} H[U] \chi \rangle_{\text{sub}} - \left(2a\mu \frac{\partial \kappa_c}{\partial \beta} + 2\kappa_c \frac{\partial(a\mu)}{\partial \beta} \right) \langle \bar{\chi} i\gamma_5 \tau^3 \chi \rangle_{\text{sub}} \right)\end{aligned}$$


- ▶ Starting point for $p(T)$ and $\epsilon(T)$ by integral method
- ▶ Subtracted expectation values: $\langle \dots \rangle_{\text{sub}} \equiv \langle \dots \rangle_{T>0} - \langle \dots \rangle_{T=0}$
→ interpolations for $T = 0$ data
- ▶ Preliminary results for $m_\pi \approx 400$ MeV and $m_\pi \approx 700$ MeV

Trace anomaly - Lattice Artifacts

$m_\pi \approx 700$ MeV:




$m_\pi \approx 400$ MeV:





Trace Anomaly, Tree Level Corrections

- ▶ Observe large lattice artifacts in $\frac{l}{T^4}$
- ▶ Lattice pressure in the free limit p_{SB}^L known: [P. Hegde *et al.*, 2008]
- ▶ Twisted mass action: [O. Philipsen & L. Zeidlewicz, 2010]
- ▶ Corrected by division by $p_{\text{SB}}^L/p_{\text{SB}}^C$ [S. Borsanyi *et al.*, 2010]

$m_\pi \approx 700$ MeV:




$m_\pi \approx 400$ MeV:




Corrected Trace Anomaly & Interpolation

$m_\pi \approx 700$ MeV:



$m_\pi \approx 400$ MeV:




Interpolation for **corrected** l/T^4 [S. Borsanyi *et al.*, 2010] :

$$\frac{l}{T^4} = \exp(-h_1 \bar{t} - h_2 \bar{t}^2) \cdot \left(h_0 + \frac{f_0 \{ \tanh f_1 \bar{t} + f_2 \}}{1 + g_1 \bar{t} + g_2 \bar{t}^2} \right)$$


Pressure and Energy Density

$$\frac{p}{T^4} - \frac{p_0}{T_0^4} = \int_{T_0}^T d\tau \frac{\epsilon - 3p}{\tau^5} \Big|_{\text{LCP}}$$

$m_\pi \approx 700$ MeV:



$m_\pi \approx 400$ MeV:



Outline

Motivation and Setup

T_c and Chiral Scenarios

Thermodynamic Equation of State

Conclusions

Conclusions & Outlook

► Conclusions:


- T_c for pion masses in the range 300 - 700 MeV
- Chiral limit so far inconclusive
- $O(4)$ scaling compatible up to scaling violation terms
- Thermodynamic equation of state for two pion masses
- Have to further improve precision at $T > 0$ and $T = 0$

► Outlook:

- $N_f = 2 + 1 + 1$

Comparison with DIK-Collaboration

(G. Schierholz et. al) from $\sigma_{\langle\bar{\psi}\psi\rangle}$:



Renormalization of $\text{Re}(L)$


$$\langle \text{Tr} \hat{L}^\dagger(\vec{x}) \text{Tr} \hat{L}(\vec{0}) \rangle = e^{-\frac{F_{\bar{q}q}(\vec{x}, T) - F_0(T)}{T}} \xrightarrow{T \rightarrow 0} e^{-\frac{V(|\vec{x}|)}{T}}$$

Renormalization of $\langle \bar{\psi}\psi \rangle$


$$\langle \bar{\psi}\psi \rangle_{ren} = Z_P \left(\langle \bar{\chi} i \gamma_5 \tau^3 \chi \rangle_{bare} + \frac{\mu c_P(\beta)}{a^2} \right) + \dots$$

$T = 0$ Subtractions & Interpolations

example: plaquette:



$\beta = 3.70$, $a\mu = 0.009$, $m_\pi \approx 400$ MeV



β -Function



[M. Cheng et al.: Phys.Rev. D77:014511, 2008]

$$\left(a \frac{d\beta}{da}\right) = - \left(\frac{r_X}{a}\right) \left(\frac{d\left(\frac{r_X}{a}\right)}{d\beta}\right)^{-1}$$

$$\left(\frac{r_X}{a}\right) (\beta) = \frac{1+n_0 R(\beta)^2}{d_0(a_{2L}(\beta)+d_1 R(\beta)^2)}$$

$$R(\beta) = \frac{a_{2L}(\beta)}{a_{2L}(3.9)}$$

$$r_0 = 0.420(15) \text{ fm}$$



Tree Level Corrections (closer look), SB (Free) Limit

Lattice:

$$\frac{p_F^L(\mu)}{T^4} = 3 \int_{[0,2\pi]^3} \frac{d^3 k}{(2\pi)^3} \frac{1}{N_t} \sum_{n=0}^{N_t-1} \ln \text{Det} (|G(k)|^2 + (a\mu)^2)$$

$$G(k) = (am) + 2r \sum_{\mu} \sin^2 \left(\frac{ak_{\mu}}{2} \right) + i \sum_{\mu} \gamma_{\mu} \sin(ak_{\mu})$$

Continuum:


$$\frac{p_F^C(\mu/T)}{T^4} = 2 \frac{7}{8} \frac{\pi^2}{90} g(\mu/T)$$
$$g(\mu/T) = \frac{360}{7\pi^4} \int_{\mu/T}^{\infty} dx \, x \sqrt{x^2 - (\mu/T)^2} \ln(1 + e^{-x})$$

Correction:

N_{τ}	4	6	8	10	12
$\rho_{\text{SB}}^L / \rho_{\text{SB}}^C$	2.586	1.634	1.265	1.134	1.084

Lines of Constant Physics, constant m_π

$m_\pi \approx 700$ MeV: presently fulfilled up to 10 %



$m_\pi \approx 400$ MeV:

