

Phase transitions and the Renormalization Group

Summer semester 2022

Problem set 1

Discussion of problems: Monday, April 28

April 21, 2022

Problem H1: Phase space and partition functions

In this problem we study phase space integrals and investigate the equivalence of the microcanonical and canonical ensemble for a free classical gas.

1. Show that the surface S_N of an N -dimensional sphere with a radius R is given by

$$S_N = \frac{2\pi^{\frac{N}{2}}}{\Gamma(\frac{N}{2})} R^{N-1} \quad (1)$$

HINT: Use the integral

$$\int d^N a e^{-a_1^2 - \dots - a_N^2} = \left(\int_{-\infty}^{\infty} da_1 e^{-a_1^2} \right)^N = \pi^{N/2}$$

and the integral representation of the Gamma function:

$$\Gamma(z) = \int_0^{\infty} x^{z-1} e^{-x} dx.$$

2. Calculate the volume of a spherical shell of thickness ΔR . Determine the fraction of the volume of this shell for $\Delta R/R = 0.01$ compared to the volume of the full sphere for $N=3, 10, 100, 500$ and 1000 dimensions.
3. Evaluate the semiclassical partition function in the microcanonical ensemble for an ideal gas in a volume V , with total energy E and N particles:

$$Z_{mc} = \int \frac{d^{3N}x d^{3N}p}{h^{3N} N!} [\Theta(E - H(p, x)) - \Theta(H(p, x) - (E - \Delta E))] \quad (2)$$

and derive the equation of state for the ideal gas:

$$PV = Nk_B T. \quad (3)$$

How do you choose the value of ΔE ?

HINT: Use the Stirling relation for large N (can you derive this relation?):

$$N! \sim N^N e^{-N}$$

4. Calculate now the canonical partition function in semiclassical approximation for an ideal gas and show that you obtain the same thermodynamic results like in the microcanonical ensemble.

Problem H2: Legendre transformation

The Legendre transformation allows to perform a variable transformation from one set of variables to a set of *canonical* variables. Consider as an example a function of two variables $f(x, y)$.

1. Consider the total derivative df . For this case there are two pairs of conjugated variables:

$$x \Leftrightarrow a \equiv \left. \frac{\partial f}{\partial x} \right|_y, \quad y \Leftrightarrow b \equiv \left. \frac{\partial f}{\partial y} \right|_x. \quad (4)$$

Show that the function $g = f - ax$ is a natural function of the variables y and a , whereas $h = f - by$ is a function of x and b .

2. Show that in classical mechanics the Hamilton function $H(q, p)$ is the Legendre transform of the Lagrange function $L(q, \dot{q})$ and the free energy $F(T, V)$ the Legendre transform of the energy E in statistical physics.
3. Consider now a function depending on N variables. How many Legendre-transforms can be in principle constructed? In which cases is the Legendre-transformed function a single-values function?
4. Consider as an example the function $f(x) = e^{x-1}$. Calculate explicitly the Legendre transform. Illustrate how to determine the Legendre-transformed function graphically. What is the interval of definition of the function?